

# Use of Performance Enhancing Drugs, Their Effects and Types Used by The Students of Faisalabad Medical University

Muhammad Ehtisham Saqib, Umme Aeman Khan, Iqra Naeem, Hafiz Yaseen

## ABSTRACT

**Objectives:** To find out the percentage of students from Faisalabad Medical University that are currently using Performance enhancing drugs, to find out the types of Performance enhancing drugs that are currently being used by the students and to find out the effects of these performance enhancing drugs on the health of the students. **Study Design:** Cross sectional study. **Setting:** 25 students each from 2<sup>nd</sup> year, 3<sup>rd</sup> year, 4<sup>th</sup> year and 5<sup>th</sup> year studying at Faisalabad Medical University were included in the study. **Study Period:** January, 2015 to June, 2015. **Materials and methods:** A total of one hundred medical students were anonymously given a 10-item questionnaire. 25 students each from 2<sup>nd</sup> year, 3<sup>rd</sup> year, 4<sup>th</sup> year and 5<sup>th</sup> year were involved in the study. The team randomly distributed the questionnaire among the students on alternative days. The completed questionnaire was sealed in envelopes and analyzed. **Results:** The age of the students completing the Performa ranged from 19-25 years. In the male population the highest use of PEDS was observed in the age group of 22 years, while the lowest use of PEDS was observed in the age group of 19 years. The highest percentage of females that used PEDS was observed in the age group of 22 years as well similar to the male ratio of PEDS use at the high level. The lowest use of PEDS was observed at the age group of 19 and 25 which was 5 % in both age groups. **Conclusion:** The results indicated that PEDS were used by the male students in a higher percentage as compared to the female students. The Caffeinated energy drinks were used more commonly. The results of this strategic survey have provided for the first-time valuable information about the use of PEDS by students of FMU to achieve desirable effects. It demonstrates that the use of PEDS in the students was high as compared to expected results; especially higher in those male students involved in sports or gym activities specifically bodybuilding athletes.

**Keywords:** Performance enhancing drugs(PEDS), anabolic steroids(AS).

### Corresponding Author

**Dr. Umme Aeman Khan**

Women Medical Officer

Govt. General Hospital, Faisalabad

Contact: +92 313-6626301

Email: drkhan185@gmail.com

*Submitted for Publication: 10-11-2017*

*Accepted for Publication: 11-02-2018*

**Article Citation:** Saqib ME, Khan UA, Naeem I, Yaseen H. Use of Performance Enhancing Drugs, Their Effects and Types Used by The Students of Faisalabad Medical University. APMC 2018;12(1):31-3.

## INTRODUCTION

Performance-enhancing Drugs (PEDS) are substances used to improve physical/athletic performance or physical appearance. The most frequently used PEDS are anabolic steroids (AS) are used primarily by males, with prevalence estimates ranging from 4% to 6%<sup>1</sup>. In addition to illegal AS, there is an abundance of legal dietary supplements available that claim to improve athletic/physical performance. Some of the supplements available replenish nutrients lost during workout and are comparatively safe (e.g. creatine), while others mimic the effects of AS (e.g., Dehydroepiandrosterone) and have similar side effects.<sup>2</sup>

The use of PEDS by teenagers has increased dramatically in the past decades. Lean mass builders, which drive or amplify the growth of muscle and lean body mass include AS that are synthetic derivatives of testosterone modified to enhance the anabolic<sup>3</sup> rather than the androgenic actions of the hormone. The anabolic effects are considered to be those promoting protein synthesis, muscle growth and erythropoiesis. There are numerous side-effects to anabolic steroids, including hypertension and atherosclerosis, blood clotting, jaundice, hepatic carcinoma, tendon damage, psychiatric and behavioral effects and, in males, reduced fertility and gynacomastia<sup>4</sup>. No

doubt, androgens exert trophic effect on skeletal and heart muscles in patients with low circulating levels of testosterone e.g pre-pubertal or hypo gonadal subjects of both genders.<sup>5</sup> Stimulants can enhance cognitive and athletic performance by increasing focus, energy, metabolic rate, and aggression, these include caffeinated energy drinks. The main ingredient of energy drinks is caffeine, along with others like taurine, pyridoxine, riboflavin, nicotinamide, vitamin B complex, and some herbal derivatives.<sup>6</sup> The acute and long-term effects results of chronic and excessive use of these drugs are not fully known. Analgesics help to perform beyond the usual nociceptive threshold. Some analgesics lead to high BP, which increases oxygen supply to muscles. Diuretics remove water from the body<sup>7</sup> and are often used by athletes who to meet weight restrictions, such as wrestlers.

The energy drink market has gained popularity rapidly. Dozens of brands are present in the market, which contain caffeine content ranging from a 50 to 505 mg per bottle<sup>8</sup> which is quite shocking. Due to regulatory oversight vigorous marketing of energy drinks, targeted specially toward young males, for performance-enhancing, psychoactive and stimulant drug effects. This has lead to increasing cases of caffeine intoxication from energy drinks, and it seems that problems like dependence

and withdrawal of caffeine will also rise. Genetics may also be vital to a person's vulnerability to caffeine intoxication, dependence, and withdrawal. The combined use of caffeine and alcohol is increasing that may increase the incidence of alcohol-related injury<sup>9</sup>. Some studies also suggest that energy drinks serve as a gateway to other forms of drug dependence.

## METHODOLOGY

**Study Design:** Cross sectional study.

**Place of Study:** Faisalabad Medical University, Faisalabad.

**Duration of Study:** January, 2015 to June, 2015.

### Methods:

A total of one hundred medical students were anonymously given a 10-item questionnaire. 25 students each from 2<sup>nd</sup> year, 3<sup>rd</sup> year, 4<sup>th</sup> year and 5<sup>th</sup> year were involved in the survey. The survey team randomly distributed the questionnaire among the students on alternative days. The completed questionnaire was sealed in envelopes and analyzed.

For all reported estimates based on the total sample the confidence level is 99% and with a margin of error of less than  $\pm 1\%$ . A survey of this nature also has some shortcomings and limitations. Inaccurate answers to the questions were an obstacle. Students asked to report their use of prohibited substances such as anabolic steroids, may be expected to underreport such use. To minimize this potential all students were guaranteed full confidentiality and asked to complete the questionnaire anonymously.

Out of the total students (n=100) 50 were males and 50 females. The survey was carried out by the cooperation of the Community medicine department, FMU. In addition, information was gathered regarding the student's knowledge and attitude regarding PEDS. Questions were also posed about the subject's behavior after taking the drugs.

A probability sampling technique was used to select a survey population of 100. The year of the students was used as an approximation of age. The study population was between 19-25 years.


Painkillers and Sedatives were also included in the survey to permit an observation of the student's perceptions and level of knowledge of a range of substances including AS and energy drinks. The students were not told whether certain substances possessed performance enhancing properties; the intention was to ascertain what they taught and believed about these substances.

## RESULTS

The age of the students completing the Performa ranged from 19-25 years. In the male population the highest use of PEDS was observed in the age group of 22 years, while the lowest use of PEDS was observed in the age group of 19 years. The highest percentage of females that used PEDS was observed in the age group of 22 years as well similar to the male ratio of PEDS use at the high level. The lowest use of PEDS was observed at the age group of 19 and 25 which was 5 % in both age groups.

**Table 1: Substances used as PEDS by percentage**

%	Caffeinated energy drinks	Painkillers	Sedatives	Anabolic steroids
overall (Mean)	24.7	13.5	10.2	2.3
19 years	25.2	7.5	10.5	2.5
20 years	28.5	8.8	9.5	1.8
21 years	25.3	12.5	12.2	3.5
22 year	28.5	14.5	13.1	2.9
23 years	23.5	12.2	9.7	2.2
24 years	21.5	19.1	7.8	1.9
25 years	20.5	20.3	9.1	1.8



**Graph 1: Percentage of drugs used by the Students by year**

## DISCUSSION

While the students were advised to complete the questionnaire anonymously, under reporting or misinformation might have been provided to some extent. The usage levels of caffeinated energy drinks was the highest among all students, which was 24.7%. 13.5 % painkillers, sedatives 10.2 % and anabolic steroids was 2.3% AS and protein supplements were used by students involved in bodybuilding or athletics to gain better sporting performances<sup>10</sup>. Educational programs regarding the use of AS must be carried out, providing information about the dangers and hazards of their continuous use. The excessive use of painkillers at such ages was quite alarming; given the stress of Medical studies and excessive exertion of their mind, headaches might be an obvious reason for their excessive use. There was no significant difference between the students who approved the use of PEDS and those who were against it. The highest percentage of drug use was found in the age range of 21 $\pm$ 2 and 22 $\pm$ 2 i-e in the students of 4<sup>th</sup> year.

The use of AS was less overall as compared to the other PEDS, but even this small percentage is alarming. This finding underscores the need for prevention programs directed at these AS users who might be termed "Body image" users. Indeed, education aimed at body image users should take into account the experiences of those who have been addressing issues related to abuse of weight-loss products and stimulants.

In the design and delivery of any interventions aimed at curbing anabolic-androgenic steroids use, it must be recognized that such behavior often produces exactly the results intended by the

user. Weight gain and change in body dimensions frequently follow AS use, and performance enhancement will occur as a consequence of the combination of AS and appropriate training regime. Because the desired effects of the drug are so evident to the user, it might be difficult to curb the user from taking such PEDS.

Educational programs aimed at decreasing the AS use must carefully address the potential health consequences, but the survey results suggest that other approaches may be beneficial. Some of the students for example believed that these AS were not harmful and were helpful in their bodybuilding process.

The high use of caffeinated energy drinks is alarming and caution should be taken in their use. These days the consumption of energy drink brands like Red Bull, Monster have become more of a fashion than a necessity<sup>11</sup>. The ingredients of these energy drinks contain high levels of stimulants, one of the most important being "taurine"<sup>12</sup>. But students feel comfortable with these drinks attributing its effects to be helpful in exam preparation for prolonged studying ability.

Painkiller use was also estimated to be quite high for such an age group of students. The high percentage of sedatives was also quite interesting, this might be regarded to the fact of stress or insomnia among the students might be a major factor of the use of these drugs.

## CONCLUSION

The results indicated that PEDS were used by the male students in a higher percentage as compared to the females specially Caffeinated energy drinks.

The results of this strategic survey have provided a reliable measure of the use of PEDS by the students to achieve desirable effects. Use of PEDS in the students was high as compared to expected results; specially in those male students involved in sports, gym and bodybuilding.

## REFERENCES

1. Advocat C. What are the cognitive effects of stimulant medications? Emphasis on adults with attention-deficit/hyperactivity disorder (ADHD). *Neurosci Biobehav Rev.* 2010;34(8):1256-66.
2. Alford C, Cox H, Wescott R. The effects of red bull energy drink on human performance and mood. *Amino Acids.* 2001;21(2):139-50.
3. Buckley WL, CE Yeasallis, KE Friedel, WA Anderson, AL Streit, JE Wright. Estimated prevalence of anabolic steroid use among high school seniors. *JAMA.* 1988;260(23):3441-5.
4. Childs E and DW Harriet, Enhanced mood and psychomotor performance by a caffeine-containing energy capsule in fatigued individuals. *Exp Clin Psychopharmacol.* 2008;16(1):13-21.
5. Celotti F, Negri Cesi P. Anabolic steroids: A review of their effects on the muscles, of their possible mechanisms of action and of their use in athletics. *J Steroid Biochem Mol Biol.* 1992;43(5):469-77.
6. Durant RH, Rickert VI, Ashwort CS, Newman C, Slavens G. Use of multiple drugs among adolescents who use anabolic steroids. *N Engl J Med.* 1993;328:922-26.
7. Dandoy C and Gereige RS. Performance enhancing drugs. *Pediatr Rev.* 2012; 33(6): 265–72.
8. Dodge TL and Jaccard JJ. The effect of high school sports participation on the use of performance substances in young adulthood. *Journal of adolescent health* 2006;39(3):367-73.
9. Edward A, Reginald S. Characteristics of steroids users in an adolescent school population. *Journal of alcohol and drug education.* 1992;38(1):43-9.
10. Faigenbaum AD, Zaichkowsky LD, Gardener DE, Micheli LJ. Anabolic steroid use by male and female middle school students. *Pediatrics.* 1998;101(5):6-7
11. Jarvis MJ. Does caffeine intake enhance absolute levels of cognitive performance? *Psychopharmacology (Berl).* 1993; 110:45-52.
12. Krowchuk DP, Anglin TM, Goodfellow DB, Stancin T, Williams P and Zimet GD. High school athletes and the use of ergogenic acid. *Am J Dis Child.* 1989;143(4):486-9.

## AUTHORSHIP AND CONTRIBUTION DECLARATION

AUTHORS	Contribution to The Paper	Signatures
<b>Dr. Muhammad Ehtisham Saqib</b> PGR Medicine, Madina Teaching Hospital, Faisalabad	Literature review, Discussion writing	
<b>Dr. Umme Aeman Khan</b> Women Medical Officer Govt. General Hospital Samanabad, Faisalabad	Research, Data collector	
<b>Dr. Iqra Naeem</b> Women Medical Officer Children Hospital, Faisalabad	Statistical Analysis	
<b>Dr. Hafiz Yaseen</b> PGR, Pediatrics Unit-1 Allied Hospital, Faisalabad	Proof reading and manuscript writing	